

Assessment:

Self-Efficacy for Negotiating Condom Use

Scale items:

1. I can ask a new partner to use condoms.
2. I can ask a partner I haven't been using condoms with to start using them.
3. I can refuse sex when I don't have a condom available.
4. I can get a partner to use condoms, even if I'm drunk or high
5. I can get every partner to use male condoms, even if they don't want to.

USE THE FOLLOWING RESPONSE CATEGORIES.

0. Cannot do at all
- 1.
- 2.
- 3.
5. Moderately certain can do
- 6.
- 7.
- 8.
- 9.
10. Certain can do

****These items were also developed by the Center for Community Health, Semel Institute-Neuropsychiatric Institute (NPI) of the University of California, Los Angeles. These are questions that have been used in various CCH assessments. Assessments will either include all of the questions or a selected portion. These questions do not represent a scale****

Response categories:

- Agree
- Mildly agree
- Mildly disagree
- Disagree

1. Condoms are easy to use.
2. Using condoms when having sex tells my partner I care about my health.
3. I am able to buy condoms.
4. I know where to get condoms.
5. I am able to carry condoms with me on a date in case I decide to have sex.
6. I know how to use a condom when I have sex with someone.
7. I am able to make sure a condom is used with a new sex partner.